

- JHARIA, H.K. and KHARE, M.N. 1986. Biological control of *Rhizoctonia bataticola* causing diseases in Soybean with *Bacillus subtilis* and *Bacillus* spp. Paper presented in the "Seminar on management of soil borne diseases of crop plants" held at Coimbatore from 8-10 Jan. (Abs.), p. 32.
- KRAFT, J.M. and PAPAIVIZAS, G.C. 1983. Use of host resistance, *Trichoderma* and fungicide to control soil borne diseases and increase seed yield of peas. *Plant Disease*, 67, 1234-1237.
- KUMAR, ANIL and GAWDA, K.T.P. 1983. Possible use of *Trichoderma harzianum* Rifai for the biological control of *Sclerotium rolfsii* Sacc. *J. Soil Biol. Ecol.*, 3, 59-61.
- RAMAKRISHNAN, G. and JEYARAJAN, R. 1986. Biological control of seedling disease of cotton caused by *Rhizoctonia solani*. Paper presented in the "seminar on Management of soil borne diseases of crop plants" held at Coimbatore from 8-10 Jan. 1986 (Abs.), p. 31.
- SINGH, P.J. and MEHROTRA, R.S. 1980. Biological control of *Rhizoctonia bataticola* on gram by coating seed with *Bacillus* and *Streptomyces* spp. and their influence on plant growth. *Plant and Soil*, 56, 475-483.

An *in vitro* Test for Evaluating the Efficacy of Mycoparasites on the Sclerotial Germination of Ergot (*Claviceps fusiformis* Lov.) of Pearl millet

L.MOHAN and R.JEYARAJAN
Department of Plant Pathology
Tamil Nadu Agricultural University
Coimbatore - 641 003

Ergot of pearl millet (*Pennisetum americanum* (L.) Leeke) caused by *Claviceps fusiformis* Loveless is causing severe losses in several high yielding varieties and hybrids of pearl millet in India. In addition to directly reducing grain yield, the admixture of sclerotia with grain poses a health hazard to consumers because of a toxic alkaloid. The methods of management of the disease include removal of sclerotia from the seed prior to planting by floatation in 10% salt solution, post harvest deep ploughing to bury sclerotia and adjustment of sowing time to avoid conditions conducive to infection. However, these methods do not give the desired level of disease control (Thakur *et al.*, 1985). Since the sclerotia constitute the primary source of inoculum, their destruction will reduce the perpetuation of the disease. Since the fungicides are not effective in this direction, antagonistic fungi like *Gliocladium virens* Miller and Foster, *Trichoderma viride* Pers and *T. harzianum* Rifai were tested to find out whether the germination of sclerotia can be delayed or inhibited by them. Kulkarni and Moniz (1974) reported that *Cerebella andropogonis* inhibited the sclerotial development of ergot of bajra. In this pathogen, Tripathi *et al.* (1981) reported that *Fusarium semibucinum* Fuckel and *Dactylium fusarioides majus* Wollen W., parasitized honeydew and sclerotial stages.

Mature sclerotia of uniform size (4.5 x 2 mm) were collected and washed with sterile water and 25 sclerotia were partially buried in sterilized sand in 9 cm Petri dish. The Petri dishes were kept at $25 \pm 2^{\circ}\text{C}$ under 12 h

darkness, alternated with light (Prakash *et al.*, 1981). The sand was kept moist by pouring sterile water at regular intervals. Two ml of the spore suspension ($10^5/\text{ml}$) of the antagonistic organisms i.e. *T. viride*, *T. harzianum* and *G. virens* was added to each Petri dish. Sterilized water was added to control plates. The treatments were replicated six times. The number of germinating sclerotia was recorded at weekly intervals for four months.

The earliest germination of sclerotia was observed after eight weeks. The control plates recorded 58.6 per cent germination (Table 1). All the three antagonistic organisms significantly reduced the germination of sclerotia and were on par with each other. The antagonistic organisms not only reduced the germination per cent but also delayed the germination of sclerotia. In the control plates, the sclerotia germinated after 9 weeks whereas none of the sclerotia in antagonist - treated plates germinated at that time. In antagonist treated plates germination took place after 12 weeks. Even after 15 weeks, the germination per cent in antagonist-treated plates was 3.3, 4.0 and 3.3 for *T. viride*, *T. harzianum* and *G. virens* respectively.

Tripathi *et al.* (1981) reported that size of sclerotia of *C. fusiformis* was reduced considerably and germinability was inhibited by 50% when they were parasitized by *F. sambucinum* and *Dactylium fusarioides*. Muller *et al.* (1985) reported that treatment of sclerotia of *Sclerotinia sclerotiorum* with *G. virens* inhibited carpogenic germination of the pathogen.

Table 1. Effect of antagonists on the germination of ergot sclerotia

Treatment	No. of sclerotia germinated (out of 150) weeks after			Per cent germination
	9	12	15	
<i>T. viride</i>	0.0	1.0	5.0	3.3 ^a
<i>T. harzianum</i>	0.0	1.0	6.0	4.0 ^a
<i>G. virens</i>	0.0	1.0	5.0	3.3 ^a
Control	66.0	82.0	88.0	58.6 ^b

In a column, means followed by similar letters are not different statistically ($P = 0.05$) by DMRT

In Maryland, USA, 100 and 1000 macro conidia of *Sporidesmium sclerotivorum* Uecker/g soil introduced in the plough layer reduced the number of sclerotia of *S.sclerotiorum* (Lib.) de Bary by 75 and 94 per cent respectively (Ayers and Adams, 1979). The results clearly indicate the potential of antagonistic fungi in reducing the inoculum load of the pathogen.

Key words : Ergot, mycoparasites, biological control

REFERENCES

- AYERS, W.A. and ADAMS, P.B. 1979. Mycoparasitism of sclerotia of *Sclerotinia* and *Sclerotium* species by *Sporidesmium sclerotivorum*. *Can. J. Microbiol.*, 25, 17-23.
- KULKARNI, C.S. and MONIZ, L. 1974. Occurrence of hyperparasite *Cerebella* on ergot of bajra. *Curr. Sci.*, 43, 803.
- MULLER, J.D., NIEDEBALKI, M., SINCLAIR, J.B. and JACOBSEN, B.J. 1985. An *in vitro* test for evaluating efficacy of mycoparasite on sclerotia of *Sclerotinia sclerotiorum*. *Pl. Dis.*, 69, 584.
- PRAKASH, H.S., SHETTY, H.S., SUBRAMANYA, S. and SAFEEULLA, K.M. 1981. Standardization of sclerotial germination technique and infectivity of ascospores of *Claviceps fusiformis* Lov. *Indian J. Agric. Sci.*, 51, 900-904.
- THAKUR, R.P., RAO, V.P., WILLIAMS, R.J., CHAHAL, S.S., MATHUR, S.B., PAWAR, W.B., NAFADE, S.D., SHETTY, H.S., SINGH, G. and BANGER, S.G. 1985. Identification of stable resistance to ergot in pearl millet. *Pl. Dis.*, 69, 982-985.
- TRIPATHI, R.K., KOLHE, S.J. and NENE, Y. 1981. Mycoparasite of *Claviceps fusiformis*, causal fungus of ergot of pearl millet. *Indian J. Mycol. Pl. Path.*, 11, 114-115.