

### ATIBALA: AN OVERVIEW

S. B. GAIKWAD\*, PROF. G. KRISHNA MOHAN

*For Author affiliations see end of the text*

This paper is available online at [www.jprhc.in](http://www.jprhc.in)

#### ABSTRACT:

Abutilon indicum is known as 'Atibala' in Sanskrit. Literally, 'Ati' means very and 'Bala' means powerful, referring to the properties of this plant as very powerful. A. indicum is a hairy herb or under shrub distributed throughout the tropica. In traditional systems of medicine, various plant parts such as roots, leaves, flowers, bark, seeds, and stems have been used as antioxidant,

#### INTRODUCTION:

A. indicum (Linn) Sweet, is a hairy herb commonly known as 'Indian mallow' belonging to family Malvaceae. It is found abundantly in the hotter parts of India but it occurs throughout the tropica, subtropica, and Ceylon<sup>[1, 2]</sup>. In addition to 'Atibala' it is also known as Thuthi, Kanghi in Hindi, and Mudra in Marathi<sup>[3]</sup>.

It grows as weed and found abundantly in wastelands from seashores 1,200 meters high in India and in sub Himalayan tracts. It is herbaceous, or shrubby, softly tomentose plant, stem is round, often tinged with purple color. The leaves are 9 by 5 cm up to petiolate, ovate to orbicular –cordate, acuminate, and toothed<sup>[4]</sup>. Flowers are borne solitary in long jointed and axillary pedicels. Calyx lobes divided in the middle and apiculate. Corolla is yellow or orange yellow opens in the evening. Carpels are 15-20 in number. Fruits are hispid, scarcely longer than the calyx and awns are erect. Seeds are 3 -5, ovoid, kidney shaped, dark brown black, tubercled or with minutely stellate hairs. Tap roots, fairly long with a number of lateral branches: 1.5-2 cm in diameter, light brown, outer surface smooth with dot like lenticels. Bark thin and can easily peel off, it has feeble odor, astringent and bitter tastes<sup>[5]</sup>.

It has a many medicinal properties and has been used by native peoples from all regions where it is found.

demulcent, laxative, diuretic, analgesic, anti-inflammatory, and antiulcer. The present review is therefore an effort to give the detailed survey of literature on its pharamcognosy, phytochemistry as well as traditional and pharmacological uses.

**Keywords:** Abutilon indicum, Atibala, Indian mallow, Pharamcognosy, Phyto-chemistry, Pharmacology.

The folk practitioner also use this plant for curing blood dysentery, fever, allergy and also aphrodisiac<sup>[6]</sup>. Bark is used in strangury and urinary complaints and is valued as a diuretic. Leaves are used for toothache, lumbago, piles and all kinds of inflammation. Decoction of leaves is used in bronchitis, catarrhal bilious diarrhoea, gonorrhoea and inflammation of bladder, and fevers. It is prescribed as a mouthwash in cases of tender gums and toothache. Seeds are tonic and Unani systems of medicine suggest its use in piles, chest troubles, bronchitis, and gonorrhoea. Rectum of children's affected with thread worms are exposed to the smoke of seeds burned on charcoal. According to the Chinese in Hong Kong, the seeds are employed as an emollient and demulcent. Infusion of root is useful in fever as a cooling medicine, strangury, heamatouria, and also in leprosy. Root is used as a pulmonary sedative and diuretic. Porter Smith states that the entire plant and seeds are used as demulcent, lenitive, diuretic, and laxative<sup>[3]</sup>.

It has been a reputed remedy in the Siddha systems of medicine for piles, jaundice, leprosy and ulcer<sup>[7]</sup>. In Vedic periods, the roots of the Bala plants i.e. Atibala (A. indicum Linn.), Mahabala (Sida rhombifolia Linn.), Bala (Sida cordifolia Linn.) and Bhumibala (Sida veronicaefolia Lam) were used to remove poison, vata – pitta diseases, heart problems, bily blood, eye diseases, and uterine disorders. Its seeds and roots both were used in

fever in the form of decoction i.e., powdered plant material dissolve in water or any other solvent <sup>[8]</sup>

### PHARMACOGNOSY:

The root of *A. indicum* is cylindrical 1.2-1.5 cm in diameter with smooth surface, having fragrance with saltish taste and yellow in color. The stem is 0.3-0.9 cm in diameter and yellow. The leaves are evergreen, stipulate, and cordate. The bark is flattened having hairy, yellow outer surface and inner surface is smooth. The fractures are fibrous. The flowers are yellow in color, pedicellate bisexual. The petiole is 1.5-7.0 cm long, yellowish brown in color, cylindrical with stellate hair. The lamina is crenate, reticulate, acute to acuminate, minutely stellate, dentate, dull green in color, hairy above and glaucous below. Glandular hairs are present while the texture is coriaceous. Fruit is a schizocarp while the seeds are furrowed, minute and glabrous <sup>[5,9]</sup>.

Microscopically the stem is undulate in outline with unicellular and multicellular hairs which are blunt. The root is also undulate in outline, with secondary wood arranged in definite rings. Some giant unicellular hairs are also present. The leaves are dorsiventral and covered with flask shaped, stellate, pitcher glandular hairs. The epidermal cells have straight anticlinal walls while the stomata are anomocytic <sup>[9]</sup>.

### PHYTO-CHEMISTRY:

The knowledge of individual chemical constituents of a medicinal plant is essential for understanding pharmacological activity as well as potential toxicity and optimizing extraction procedures.

#### Leaves:

Leaves contain tannins, mucilage, traces of asparagin, organic acid and, ash of leaves contains alkaline sulphates, chlorides, magnesium phosphate and calcium carbonate <sup>[1]</sup>. Ethanolic extract contain 72% more quercetin than flowers <sup>[10]</sup>. Leaves also contains alkaloids, sterols, terpenoids, glycosides <sup>[5]</sup> essential oils as well various amino acids <sup>[11]</sup>. Baxi et al isolated tocopherols and  $\beta$  – sitosterol from leaves <sup>[12]</sup>.

#### Aerial parts

The aerial part of the plant on extraction with petroleum ether led to the isolation of n-alkane mixture, an alkanol fraction and  $\beta$  – sitosterol; fumaric, p-coumaric, vanillic, caffeic, and p-hydroxybenzoic, p -  $\beta$  -D-glucosyloxybenzoic acids, and gluco-vanilloyl glucose, fructose, aspartic acid, histidine, threonine, serine, and leucine. Galactose and galacturonic acids are present in mucilage fraction. Saponins, flavnoids, and alkaloids are present in shoot and flowers <sup>[13]</sup>.

#### Roots

Roots contain asparagin <sup>[14]</sup>. Gallic acid <sup>[15]</sup> and Fixed oil were isolated from root <sup>[16]</sup>. Bhattacharjee reported presence of sterols, terpenoids, terpens, flavonoids, and steroids <sup>[17]</sup>.

#### Flowers

Gossypetin-8 and 7-glucosides and cyanidin-3-rutinoside isolated from flower petals of *A. indicum* <sup>[18]</sup>. Two sesquiterpene lactones i.e alantolactone and isoalantolactone have been first time reported by Sharma et al <sup>[19]</sup>. Some flavonoids like luteolin, chrysoeriol, luteolin 7-O- $\beta$  -glucopyranoside, chrysoeriol 7-O- $\beta$  -glucopyranoside, apigenin 7-O- $\beta$  -glucopyranoside, quercetin 3-O- $\beta$  -glucopyranoside, quercetin 3-O- $\alpha$ -rhamnopyranosyl (1 - 6)- $\beta$ -glucopyranoside, were isolated and identified <sup>[20]</sup>. Oil obtained from the flowering tops yielded geraniol, geraniol actate,  $\alpha$ - pinene, borneol, and tetradecane <sup>[21]</sup>.

#### Seeds

Seeds contain water soluble galactomannan contains D-galactose and D- mannose in 2:3 ratios. Acid catalysed fragmentation, periodate oxidation and methylation showed that the seed-gum has branched structure consisting of linear chain  $\beta$ -D (1 4) linked mannopyranosyl units, some of which are substituted at ortho-6 by two  $\alpha$  -D (1 6) galactopyranosyl units mutually linked glycosidically as end groups <sup>[22]</sup>. Chemical analysis of the seed oil showed the presence of stearic, linolenic, oleic, and palmitic acid <sup>[23]</sup>. Seeds were analyzed for the crude pentosan, protein, and water soluble mucilage

contents<sup>[24]</sup>. HBr reactive fatty acids viz, 12, 13-epoxyoleic (vernolic acid); 9, 10 methylene-heptadec-8-enoic (Malvalic acid) were identified in the seed oil<sup>[25]</sup>. Amino acid profile of seed proteins (31%) contains threonine, glycine, serine, glutamine, lysine, methionine, isoleucine, proline, alanine, cysteine, tyrosine, phenylalanine, leucine, asparagine, histidine, valine, arginine<sup>[26]</sup>. TLC-GLC studies of seed oil revealed the presence of high amount of unsaturated acids. Stearic acid and palmitic acid were the principal component from the saturated acids. Raffinose as a prime sugar component was found in seed<sup>[27]</sup>.

### Fruits

Fruits contain flavanoids and alkaloids<sup>[28]</sup>.

### Whole plant:

Some flavonoids have been isolated from *A. indicum* like quercetin, kaempferol, gossypetin, and cyanidin glycosides<sup>[29]</sup>. The investigation on the chemical constituents of the whole plant has resulted in the isolation of two new compounds, abutilin A (1) and (R)-N-(1'-methoxycarbonyl-2'-phenylethyl)-4-hydroxybenzamide (2), as well as 28 known compounds<sup>[30]</sup>.  $\beta$ -sitosterol as a potential new mosquito larvicidal compound was isolated from petroleum ether extract<sup>[31]</sup>. The plant was found to contain gum resin and mucilage<sup>[32]</sup>. Tannins were not present in 50% ethanolic extract of the plant<sup>[33]</sup>.

The essential oil of plant contains  $\beta$ -pinene, caryophyllene, caryophyllene oxide, 1, 8-cineole, ceraneol, ceranyl acetate, elemenes, eudesmol, and farnesol<sup>[34]</sup>. Preliminary phytochemical test shows the presence of glycosids, leucoanthocyanidin, saponins alkaloids cardiac glycosides, cyanogenetic tannins, and, phenolic compounds in the leaves, root, and stem<sup>[9]</sup>.

### PHARMACOLOGY:

#### a) Antioxidant and radical scavenging activity:

The roots and aerial parts were extracted in n-hexane, chloroform, butanol, and ethyl acetate, evaluated for their total phenolic content, total flavonoid content, total antioxidant capacity (TAC) and Trolox equivalent antioxidant capacity (TEAC), employing 2,2'-azinobis(3-ethylbenzothiazol-6-sulfonic acid) (ABTS) and ferric

reducing anti-oxidant power (FRAP). TEAC values ranged from 3.019 to 10.5  $\mu$ M for n-hexane and butanol fractions using the ABTS assay. The FRAP assay showed reducing powers of the fractions in the order of butanol > ethyl acetate > chloroform > n-hexane and butanol > chloroform > hexane > ethyl acetate. By using the DPPH free radical assay  $T_{(EC50)}$  and  $EC_{50}$  values were determined. The antioxidant/radical scavenging capacity of the extracts was found to be a dose-dependent activity. From this report one can conclude that plant is a potent source of natural antioxidant<sup>[35]</sup>.

#### b) Antidiabetic activity:

The aqueous extract was administered to moderately diabetic rats at a dose of 0.5 and 1 g/kg body weight in an oral glucose tolerance test led to a significant reduction in plasma glucose levels in 30 minutes as compared with untreated rats ( $P < 0.05$ ), and it was at a faster rate than the use of standard antidiabetic drug, i.e. glibenclamide. By using an everted intestinal sac inhibition of glucose absorption through the small intestine was investigated. The extract at concentrations of 0.156 to 5 mg/mL caused a reduction of glucose absorption in a dose response manner. The maximum response was noted at a dose of 2.5 mg/mL. From above observations one can predict that the aqueous extract of the *A. indicum* plant has good antidiabetic property<sup>[36]</sup>.

#### c) Antimalarial activity

Larvicidal activity of crude hexane, petroleum ether, ethyl acetate, acetone and methanol extracts of *Abutilon indicum*, were tested for their toxicity against the early fourth-instar larvae of *Culex quinquefasciatus*. All extracts showed moderate larvicidal effects after 24 h. Petroleum ether extract showed highest larval mortality.  $\beta$ -sitosterol isolated from plant was a potential new mosquito larvicidal compound with LC50 value of 11.49, 3.58 and 26.67 ppm against *Aedes aegypti* L, *Anopheles stephensi* Liston and *Culex quinquefasciatus* Say (Diptera: Culicidae), respectively<sup>[31]</sup>.

#### d) Hepatoprotective activity

The aqueous leaf extract of *A. indicum* exhibited significant hepatoprotective activity in carbon tetrachloride- and paracetamol-induced hepatotoxicities in rats by reducing carbon tetrachloride- and paracetamol-

induced change in bio-chemical parameters like SGPT, SGOT, and ALKP by enzymatic examination. Hepatoprotective action was found may be due to interference of leaf extract with free-radical formation. Aqueous extract at a dose of 100 and 200 mg/kg body wt p.o showed hepatoprotective activity comparable to standard drug Silymarin (100 mg/kg body wt). LD50 value is more than the dose of 4 g/kg body wt in acute toxicity models [37].

### e) Hypoglycemic activity

Alcohol and water extracts of *A. indicum* leaves at a dose of 400 mg/kg, p.o showed significant hypoglycemic effect in normal rats 4 hrs after administration (23.10% and 26.95%, respectively). When results were compared to standard drug Tolbutamide alcohol and water extracts showed weak activity while petroleum ether and chloroform extract did not showed significant hypoglycemic activity [6].

### f) Analgesic activity

Isolated eugenol [4-allyl-2-methoxyphenol], from *A. indicum* was found to possess significant analgesic activity. At doses of 10, 30, and 50 mg/kg body weight, it exhibited 21.30 ( $p < 0.05$ ), 42.25 ( $p < 0.01$ ) and 92.96% ( $p < 0.001$ ) inhibition of acetic acid induced writhing in mice. In the radiant heat method at a dose of 50 mg/kg body weight eugenol showed 33.40% ( $p < 0.05$ ) prolongation of tail flicking time [38].

Fixed oil isolated from petroleum ether extract of the root showed dose –dependent analgesic activity in rats using analgesiometer and in mice against acetic acid induced writhing test. Fixed oil exhibited analgesic effect at a dose of 400mg/kg s.c. LD50 of the fixed oil in mice was 2357.9 mg/kg p.o and 933.3 mg/kg s.c. No behavioral changes were observed in mice [16, 39]. Gallic acid in rat showed analgesic activity at dose of 0.1gm/kg i.p within 30–45 minutes of administration [15]. Central analgesic activity of fixed oil isolated from *A. indicum* tested on animal at doses of 400 and 600 mg/kg i.p by using tail flick response to radiant heat, acetic acid induced writhing, and tail clip method [40].

### g) Estrogenic/antiestrogenic properties

Methanolic extracts of fruits at a dose of 100-500 mg/kg body weight caused suppression of uterine peroxidase enzyme activity. It was suggested that the uterine peroxidase assay could be utilized as a biochemical parameter in the screening of new antifertility agents for their estrogenic/antiestrogenic properties [41].

### h) Immunomodulatory activity:

Essential oil from plant augments antibody in animals showing immunological value [11]. *A. indicum* was evaluated experimentally for its immunostimulant properties. The parameters included i) humoral antibody enhancing response of drug against killed salmonella typhi ‘O’ antigen in rabbits, ii) protective effect of drugs against *Staphylococcus aureus* (Coagulase positive) challenge in rabbits iii) direct antibacterial activity of drugs against *Staphylococcus aureus* on nutrient agar plate and iv) immunological changes in experimental animals following drug administration. Experimental studies in rabbits showed that the entire drug treated groups of animals showed a higher level of anti-salmonella typhi ‘O’ titers. The values of titers were statistically significant for 8<sup>th</sup>, 16<sup>th</sup>, and 24<sup>th</sup> day, respectively when compared to the control group. Higher level of antibody titre was obtained on 24<sup>th</sup> day of experiment. The protective effect of *A. indicum* against virulent *staphylococcus aureus* challenge in the rabbits showed highly significant survival period ( $p < 0.01$ ). The haematological values for total and differential leucocytes count and Haemoglobin percent did not shown any particular change. However a significant decrease in Erythrocyte Sedimentation Rate levels ( $p < 0.05$ ) following *Staphylococcus aureus* injection was noted in *A. indicum* treated animals. Histopathological studies of animals challenged with virulent *staphylococcus aureus* showed marked protection of tissue damage [42]. The use of *A. indicum* as an adjuvant in immunization programme has been advocated due to its immunomodulatory properties [43]. The whole fine powder of the plant at a dose of 500 mg/kg body weight, when compared to the control group showed statistically highly significant rise in modulatory behaviour in all the models [44].

### **i) Antimicrobial activity:**

Essential oil obtained from whole plant exhibited antibacterial activity against *Proteus vulgaris*, *Bacillus subtilis*, *Bacillus anthracis* and while it was inactive against *Salmonella typhimurium*, *Salmonella pullorum*, and *Klebsiella* Species<sup>[45]</sup>. The aqueous extract of the whole plant at a concentration of 5-10 mg/ml showed in vitro antibacterial activity against *E. Coli*, *Corynebacterium diphtheriae*, *Streptococcus viridans*, *Salmonella typhi*, *Shigella flexneri*, *Diplococcus pneumoniae*, *Salmonella paratyphi A* and *B*. The aqueous extract of plant was devoid of antimicrobial activity against *Trichosporon cutaneum*, *Candida albicans*, *Candida tropicalis*, *Microsporium canis*, *Microsporium nanum*, *Microsporium gypseum*, *Piedraria hortae*, *Phialophora jeanselmei*, *Trichophyton mentagrophytes*, *Trichophyton rubrum*, *Madurella mycetomy*, *Histoplasma capsulatum*, and *Cryptococcus neoformans*<sup>[46]</sup>.

The 80% ethanolic extract of root was found to be devoid of any activity against *Pseudomonas aeruginosa*, *Escherichia Coli*, *Staphylococcus aureus*, and *Klebsiella aerogenes*. Benzene extract was active against *Pseudomonas aeruginosa*, *Escherichia Coli*, and *Proteus species*<sup>[47]</sup>. The hexane extract was active against *Staphylococcus aureus*, and *Pseudomonas aeruginosa*. The chloroform and ethyl acetate extract showed good activity against most of bacteria. The hexane extract showed activity against the fungi *Aspergillus terreus*, *Aspergillus ochraceus*, *Aspergillus flavus*, *Aspergillus cardius*, and *Aspergillus oryzae*. The methanolic extract was active against *Aspergillus oryzae*, *Aspergillus ochraceus*<sup>[48]</sup>.

The aqueous residue of the leaves were devoid of any antibacterial activity against *Aeromonas hydrophilla*, *Alkaligenes viscolactis*, *Staphylococcus pyrogenes*, *Escherichia Coli*, *Cytophaga species*, *Klebsiella aerogenes*, *Vibrio parahaemolytica*, *Pseudomonas aeruginosa*, *Vibrio damsela*, *Streptococcus pyrogens*, and *Bacillus cereus*<sup>[49]</sup>. The 50% acetone extract of the seeds showed antifungal activity against the plant fungi *Microsporium gypseum* (5%) and *Epidermophyton floccosum* (20%) but was found to be inactive against *Paecilomyces mentafrophytus*<sup>[50]</sup>.

Crude ethanol extract of *A. indicum* exhibits antimicrobial activity against the Gram negative organisms *P. Aeruginosa*, *P. Vulgaris*, *E. Coli*, *S. Sonne* and Gram positive organisms such as *B. Subtilis*, *B. Megatherium S. Leuka*, *S. Aureus* with a maximum diameter of zone of inhibition ranging from 24.3 mm, 22.4 mm, 20.1 mm, 24.5 mm, 22.4 mm, 21.0 mm, 23.5 mm and 24.1 mm respectively in the agar-well diffusion method<sup>[51]</sup>. The ethanolic extract of the root was active against the fungi *Aspergillus flavus*, *Aspergillus glaucus*, *Aspergillus ochraceus*, *Aspergillus oryzae*, and *Aspergillus cardius*, *Penicillium Species*. Acetone was active extract against *Aspergillus ochraceus*<sup>[52]</sup>.

### **j) Angiotensin Converting Enzyme (ACE) inhibitory activity**

The root extract of *A. indicum* was found to inhibit ACE in water, acetone, ethanol (96%), by 18 %, 9 % 1 %, and respectively. Inhibition was measured from the enzymatic cleavage of the chromophore- fluorophore-labelled substrate dansyltriglycine into dansylglycine and diglycine. It reflects the use of plant in hypertension<sup>[53, 54]</sup>.

### **k) Anti-inflammatory activity**

Ethanol, chloroform, and aqueous extract (1, 5, 10, 20 mg/100ml) of leaves of *A. indicum* were screened for anti-inflammatory activity. Prevention of hypotonicity induced HRBC membrane lysis was taken as a measure of anti-inflammatory activity. All the extracts showed biphasic effect on HRBC membrane stabilization and have comparable activity to that of Diclofenac<sup>[55]</sup>.

### **l) Miscellaneous:**

Fixed oil showed depressant action on isolated frog's heart, and had no effect on isolated guinea pig ileum. It inhibits acetylcholine- induced contraction of isolated guinea pig ileum<sup>[39]</sup>. Fruit decoction mixed with ammonium chloride is given orally to treat hemorrhagic septicemia<sup>[51]</sup>. Whole plant extract of *A. indicum* is devoid of insecticidal properties<sup>[33]</sup>. Methanolic extract (0.1 mg/ml) of whole plant was screened for acetylcholinesterase inhibitor activity using Ellman's

colorimetric method in 96-welled micro plate. Extract showed 30.66% inhibitory activity. Acetylcholinesterase inhibitors have therapeutic applications in the treatment of Alzheimer's disease, senile dementia, myasthenia gravis, ataxia, and Parkinson's disease [56]. Ethanolic extracts of plant against a P<sub>388</sub> Lymphocytic leukemia in mice in a preliminary biological screening showed anticancer activity [52]. Whole plant was screened for in-vitro anti-arthritic activity. The effect was observed on inhibition of protein denaturation, proteinase inhibition, and membrane stabilization at a dose of 100 and 200 mcg /ml. It showed significant protection against hypotonic saline induced RBC membrane damage, denaturation of proteins and possesses good anti-proteinase activity [57]. The leaf juice when mixed with jaggery is used for the treatment of snakebite as antidote [58]. The fruit is used to treat piles, gonorrhoea and cough [59, 60].

### Uses in Commercial Formulations

The plant has many Ayurvedic and commercial applications. The commercially available products are Atibalaghrit, Mahanarayan taila, Bala taila, and Mahavishgarbh taila [61, 62]. It also forms one of the ingredients of Chayvanprash Linctus, used as a general tonic for restoring strength and health and Garbha Cintamani Rasa, a classical Ayurvedic preparation which is used in puerperal complications [63].

### CONCLUSION

The extensive survey of literature revealed that *A. indicum* is an important plant of Ayurveda. It is being used since a long time in making ayurvedic drugs. Medicinally saponins possess hypoglycaemic and antifungal activities. Linoleic, oleic, palmitic, lauric, stearic, and other fatty acids found in the plant claims analgesic activities.  $\beta$ -sitosterol is reported to possess antipyretic actions and flavonoids having hypoglycaemic activities. Besides, the gum and resin obtained from the plant are used in rheumatism and show antiplasmitary reaction, which support the use of this plant for the various purposes since ancient times. Thus, the phytochemicals found therein possess various applications in the allopathic treatment for better drug rationale therapy.

### REFERENCES

1. Panda H, Herbs Cultivation and Medicinal Uses, National Institute of Industrial Research, Delhi, 2000, 98-99.
2. Chopra RN, Nair SL, Chopra IC, Glossary of Indian Medicinal Plants, CSIR, New Delhi, 1956, 2.
3. Kirtikar KR, Basu BD, Indian Medicinal Plants, Vol I, 2<sup>nd</sup> Ed, Popular Prakashan, Bombay, 1991, 314-316.
4. Sivarajan VV, Ayurvedic Drugs and Their Plant Sources, Indira Balachandran Oxford and IBH Publishing Co Pvt Ltd, New Delhi, 2004, 78.
5. The Ayurvedic Pharmacopoeia of India, Vol-I, 1<sup>st</sup> Ed, Government of India, Ministry of Health and Family welfare, Department of Indian System of Medicine and Homoeopathy, The controller of Publication, Civil Lines, New Delhi, Part-I, 1990, 20-21.
6. Seetharam YN, Chalageri G, Setty SR, Bheemachar, Hypoglycemic activity of *Abutilon indicum* leaf extracts in rats, *Fitoterapia*, Vol 73, 2002, 156-159.
7. Yoganarasimhan SN, Medicinal Plants of India, Vol 2, Cyber Media, Bangalore, 2002, 10.
8. Dwivedi KD, Vedo Mein Ayurved, Vishwabharti Anusandhan Parishad, Gyanpur (UP). 2000, 259-285.
9. Dhanalakshmi S, Lakshamanan KK, Subramania MS, Pharmacognostical and Pharmacological studies on two species of *Abutilon L*, *Journal of Research Education Indian Medicine*, Vol 9, 1990, 21-32.
10. Rajalakshmi PV, Kalaiselvi Senthil K, Direct HPLC analysis of quercetin in exudates of *Abutilon Indicum* (Linn). *Malvaceae*, *Journal of Pharmaceutical Science and Technology*, Vol. 1 (2), 2009, 80-83.
11. Database on Medicinal Plants used in Ayurveda, Vol I, Central Council for Research in Ayurveda & Sidha, New Delhi, 2002, 50.
12. Baxi AJ, Parikh AR, Isolation of some non saponifiable principle from the leaves of *A. indicum* G Don, *Bull Med Ethenobotany Res*, Vol 1, 1980, 534-537.

13. Gaind KN, Chopra KS, Phytochemical investigations of *Abutilon indicum*, *Planta Med*, Vol 30, 1976, 174-185.
14. Nadakarni KM, Nadakarni AK, *Indian Materia Medica*, Vol I, Popular Prakashan, Bombay, 1976, 8-9.
15. Sharma PV, Ahmad ZA, and Sharma VV, Analgesic constituent of *Abutilon indicum*. *Indian Drugs*, Vol 26, 1989, 333.
16. Bagi MK, Kalyani GA, Dennis TJ, Kumar AK, Kakarani HK, A preliminary phytochemical screening of *Abutilon indicum*: I. analgesic activity, *Indian Drugs*, Vol 22, 1984, 69-72.
17. Bhattacharjee AK, Das AK, Phytochemical Screening of some Medicinal plants. *Quart J Crude Drug Res*, Vol 9, 1969, 1408-1412.
18. Subramanian SS, Nair AGR, Flavanoids of four Malvaceous plants, *Phytochemistry*, Vol 11, 1972, 1518-1519.
19. Sharma PV, Ahmad ZA, Two sesquiterpene lactones from *Abutilon indicum*, *Phytochemistry*, Vol 28, 1989, 3525.
20. Matlawska I, Sikorska M, Flavonoid compounds in the flowers of *Abutilon indicum* (L.) Sweet (Malvaceae), *Acta Pol Pharm*, Vol 59(3), 2002, 227-229.
21. Geda A, Gupta AK, Chemical investigation of essential oil of *Abutilon indicum*, *Perfum Flav*, Vol 18, 1983, 39.
22. Singh V, Mishra UC, Khare GC, Gupta PC, A neutral seed gum from *Abutilon indicum*, *Carbohydrate Polymers*, Vol 33, 1997, 203-205.
23. Gambhir IR, Joshi SS, Chemical investigations of seeds of *Abutilon indicum* G. Don, *Journal of Indian Chemical Society*, Vol 29, 1952, 451-454.
24. Kapoor VP, Khan PSH, Raina RM, Farooqui MIH, Chemical analysis of seeds: Part III From 40 non- leguminous species, *Science Culture*, Vol 41, 1975, 336-340.
25. Babu M, Husain S, Ahmad MU, Osman SM, *Abutilon indicum* seed oil- Characterisation of HBr-reactive acids, *Fette Siefen Anstrichm*, Vol 82, 1980, 63-66.
26. Prakash D, Jain RK, Misra PS, Amino acid profiles of some under-utilised seeds, *Plant Food Human Nutrition*, Vol 38, 1988, 235-241.
27. Badami RC, Deshpande GS, Shanbhag MR, Minor seed oils, VII. Examination of seed oils by gas- liquid chromatography, *Journal of oil Technology assoc India*, Vol 7, 1975, 76-77.
28. Sinha SP, Dogra JVV, A survey of the plants of Bhagalpur and Santhal Pargana for saponin, flavonoids and alkaloids, *Int Journal of Crude Drug Research*, Vol 23, 1985, 77-86.
29. Phytochemical reports, *Phytochemistry*, Vol 11, 1972, 1491-1492.
30. Kuo PC, Yang ML, Wu PL, Shih HN, Thang TD, Dung NX, Wu TS, Chemical constituents from *Abutilon indicum*, *Journal of Asian Natural Products Research*, Vol 10 (7-8), 2008, 699-703.
31. Abdul Rahuman A, Gopalakrishnan G, Venkatesan P, Geetha K, Isolation and identification of mosquito larvicidal compound from *Abutilon indicum* (Linn.) Sweet, *Parasitology Research*, Vol 102 (5), 2008, 981-988.
32. Yoganarasisimhan SN, Nair KV, Keshavamurthy KR, *Medico-botany of Karnataka-3. Utilisation of floristic wealth for the economic development of Kanakapura, Taluka, Bangalore district*, *Journal of Economy, Taxonomy and Botany*, Vol 6, 1985, 97-108.
33. Atal CK, Srivastava JB, Wali BK, Chakravarty RB, Dhawan BN, Rastogi RP, Screening of Indian Medicinal Plants for biological activity Part VIII, *Indian Journal of Experimental Biology*, Vol 16, 1978, 330-349.
34. Jain PK, Sharma TC, Bokadia MM, Chemical investigation of essential oil of *Abutilon indicum*, *Acta Cienc Indica*, Vol 8C, 1982, 136-138.
35. Yasmin S, Kashmiri MA, Asghar MN, Ahmad M, Mohy-Ud-Din A, Antioxidant potential and radical scavenging effects of various extracts from *Abutilon indicum* and *Abutilon muticum*, *Pharmaceutical Biology*, Vol 48(3), 2010, 282-289.
36. Krisanapun C, Peungvicha P, Temsiririrkkul R, Wongkrajang Y, Aqueous extract of *Abutilon indicum* Sweet inhibits glucose absorption and stimulates insulin secretion in rodents, *Nutrition Research*, Vol 29(8), 2009, 579-587.

37. Porchezian E, Ansari SH, Hepatoprotective activity of *Abutilon indicum* on experimental liver damage in rats, *Phytomedicine*, Vol 12(1-2), 2005, 62-64.
38. Ahmed M, Amin S, Islam M, Takahashi M, Okuyama E, Hossain CF, Analgesic principle from *Abutilon indicum*, *Pharmazie*, Vol 55(4), 2000, 314-316.
39. Bagi MK, Kalyani GA, Dennis TJ, Kumar AK, Kakarani HK, A preliminary phytochemical screening of *Abutilon indicum*: II Analgesic activity, *Fitoterapia*, Vol 56, 1985, 169-172.
40. Almeida RN, Navarro DS, Barbosa-Filho JM, Plants with central analgesic activity, *Phytomedicine*, Vol 8(4), 2001 310-322.
41. Johri RK, Pahwa GS, Sharma SC, Zutshi U, Determination of estrogenic/antiestrogenic potential of antifertility substances using rat uterine peroxidase assay, *Contraception*, Vol 44(5), 1991, 549-557.
42. Dixit SP, Tewari PV, Gupta RM, Experimental studies on the immunological aspects of *Atibala* (*A. indicum* Linn Sw.), *Mahabala* (*Sida rhombifolia* Linn.), *Bala* (*Sida cordifolia* Linn.) and *Bhumibala* (*Sida veronicaefolia* Lam). *J Res Indian Med Yoga Homeopathy*, Vol 13 (3), 1978, 50-66.
43. Tewari PV, Sharma RD, Immunization through Ayurvedic drugs, *J Res Educ Indian Med*, Vol 11(4), 1992, 1-5.
44. Singh D, Gupta SR, Modulatory influence of *Abutilon indicum* leaves on Hepatic Antioxidant status and Lipid Peroxidation against alcohol induced liver damage in Rats, *Pharmacology online*, Vol 1, 2008, 253-262.
45. Geda A, Jain PK, Bokardia MM, Antimicrobial activity of some Indian essential Oils, *Acta Cienc Indica*, Vol 4, 1978, 248-251.
46. Naqvi SAH, Khan MSY, Vohora SB, Antibacterial, antifungal, and anthelmintic investigations on Indian Medicinal Plants, *Fitoterapia*, Vol 62, 1991, 221-228.
47. Valsaraj R, Pushpangadan P, Smitt UW, Adsersen A, Nyman U, Antimicrobial screening of selected medicinal plants from India, *Journal of Ethnopharmacology*, Vol 58, 1997, 75-83.
48. Mehta BK, Neogi R, Kotra S, Mall OP, Antimicrobial activity of *Abutilon indicum*, *Fitoterapia*, Vol 68, 1997, 273-274.
49. Srinivasan D, Nathan S, Suresh T, Perumalsamy L, Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine, *Journal of Ethnopharmacology*, Vol 74, 2001, 217-220.
50. Rathore A, Misra N, Antifungal properties of some plant extracts, *J. Science Research*, Vol 9, 1987, 103-107.
51. Prabahar AE, Thangabalan B, Chavala AS, Sasi Kumar, Kathiravan M, Karthikeyan R, Antibacterial activity of various extracts of *Abutilon indicum* (L.) Sweet leaves, *Journal of Pharmacy Research*, Vol 2(8), 2009, 1324-1325.
52. Bhakuni DS, Dhar ML, Dhar MM, Dhawan MN, Gupta B, Srimal RC, Screening of Indian Medicinal Plants for biological activity Part III, *Indian Journal of Experimental Biology*, Vol 9, 1971, 91-102.
53. Ali ZA, Folk Veterinary Medicines in Moradabad District (Uttar Pradesh), India. *Fitoterapia*, Vol 70, 1999, 340-347.
54. Hansen K, Nyman U, Smitt UW, Adsersen A, Gudiksen L, Rajasekharan S, Pushpangadan P, In vitro screening of traditional medicines for anti-hypertensive effect based on inhibition of the angiotensin converting enzyme (ACE), *Journal of Ethnopharmacology*, Vol 48, 1995, 43-45.
55. Rajurkar R, Jain R, Mataka N, Aswar P, Khadbadi SS, Anti-inflammatory action of *Abutilon indicum* (L.) Sweet leaves by HRBC membrane stabilization, *Research journal of Pharmaceutics and technology*, Vol 2(2), 2009, 415-416.
56. Mukherjee PK, Kumar V, Mal M, Houghton PJ, Acetylcholinesterase inhibitors from plants, *Phytomedicine*, Vol 14, 2007, 289-300
57. Deshpande V, Jadhav VM, Kadam VJ, In-vitro anti-arthritis activity of *Abutilon indicum* (Linn.) Sweet, *Journal of Pharmacy Research* Vol 2(4), 2009, 644-645.
58. Mohapatra SP, Sahoo HP, An Ethno-Medico-Botanical Study of Bolangir, Orissa, India: Native Plant Remedies against Gynecological



- Diseases, Ethno botanical Leaflets, Vol 12, 2008, 846-850.
59. Samy PR, Thwin MM, Gopalakrishnakone P, Ignacimuthu S, Ethnobotanical Survey of folk plants for the treatment of snakebites in Southern part of Tamilnadu, India. Journal of Ethnopharmacology, Vol 115, 2008, 302-312.
60. Ignacimuthu S, Ayyanar M, Sakarasivaraman K. Ethnobotanical study of medicinal plants used by Paliyar tribals in Theni district of Tamil Nadu, India, Fitoterapia, 79, 2008, 562-568.
61. Chatterjee A, Pakrashi SC, The Treatise on Indian Medicinal Plants, Vol 2, Publication and Information Directorate, CSIR, New Delhi, 1992, 174-175.
62. Dhiman AK, Ayurvedic Drug Plants, Daya Publishing House, New Delhi, 2006, 58-60.
63. Bulbul IJ, Ullah MO, Rahman M, Rahman KA, Paul AK, Choudhuri MSK, Effect of “Garbha Cintamani Rasa”, an Ayurvedic Formulation on Lipid Profile, Liver Function and Kidney Function Parameters of Rat Plasma after Chronic Administration, European Journal of Scientific Research, Vol 32 (1), 2009, 25-32.

### AUTHORS AFFILIATIONS AND ADDRESS FOR CORRESPONDENCE:

SWITI GAIKWAD, Research scholar,

Center for Pharmaceutical Sciences, Institute of Science and Technology, Jawaharlal Nehru Technological University, Kukatpally, Hyderabad - 500 085 Andhra Pradesh, India. (+91) 9298705073  
gaikwad\_sweety@rediffmail.com

PROF. G. KRISHNA MOHAN

Head of the Department, Center for Pharmaceutical Sciences, Institute of Science and Technology, Jawaharlal Nehru Technological University, Kukatpally, Hyderabad - 500 085 Andhra Pradesh, India. (+9140) 23158666

krishna\_mohan56@yahoo.co.in